

AstraZeneca 

# Patient Safety Information Card

**SOLIRIS<sup>®</sup> (eculizumab)**

Patients receiving SOLIRIS<sup>®</sup> should carry this card with them at all times.  
Show this card to any healthcare professional involved in your or the child's treatment.



## Important Safety Information for Patients

SOLIRIS® can lower the ability of your immune system to fight infections, especially meningococcal infection, which requires immediate medical attention. If you experience any of the following symptoms, you should immediately call your doctor or seek emergency medical care, preferably in a major emergency medical care center.



### For Pediatric Patients

Fill out the card and give one to anyone responsible for the care of your child. Also, be sure to carry a copy with you at all times. Tell the person responsible for the care of your child that these cards should be shown to any healthcare professional involved in your child's treatment, should your child need medical attention.

If your child shows any signs or symptoms of meningitis or severe blood infection (sepsis), contact their healthcare professional immediately. If you cannot reach the healthcare professional, seek immediate emergency care at an emergency department and show the staff the child's Patient Safety Information Card.



### For Supervising Individuals of Pediatric Patients

This child is currently being treated with SOLIRIS® and may have reduced natural resistance to infections, especially meningococcal infections, which include meningitis and severe blood infection or blood poisoning (also known as sepsis). If you notice that the child has any of the signs or symptoms listed on this card, which could indicate a serious infection, call the child's doctor or seek immediate emergency care at an emergency department and show this card to them.

## Common signs and symptoms of meningitis and severe blood infection

- Headache with nausea or vomiting
- Headache with a stiff neck or stiff back
- Confusion
- Muscle aches with flu-like symptoms
- Eyes sensitive to light
- Fever
- Rash

## Common signs and symptoms of meningitis and severe blood infection in infants and children

- Fever, cold hands and feet
- Fretful, dislike being handled
- Rapid breathing or grunting
- Unusual cry, moaning
- Stiff neck, dislike bright lights
- Refusing food and vomiting
- Drowsy, floppy, unresponsive
- Pale, blotchy skin spots/rash
- Tense, bulging fontanelle (soft spot on the baby's head)
- Convulsions/seizures

### In children, additional signs and symptoms may include:

- Severe muscle pain
- Severe headache
- Confusion
- Irritability



**Get emergency medical care right away if you have any of these signs or symptoms and show this card.**

**Even if you stop using SOLIRIS<sup>®</sup>, keep this card with you for 3 months after your last SOLIRIS<sup>®</sup> dose. Your risk of meningococcal infection may continue for several weeks after your last dose of SOLIRIS<sup>®</sup>.**



# Information for the Treating Physician



This patient has been prescribed SOLIRIS® (eculizumab) therapy, which increases the patient's susceptibility to meningococcal infection (*Neisseria meningitidis*) or other general infections.

- Meningococcal infections may become rapidly life-threatening or fatal if not recognised and treated early.
- **Evaluate immediately if infection is suspected and treat with appropriate antibiotics if necessary.**
- Contact prescribing physician (refer to below) as soon as possible.

For more information about SOLIRIS®, please refer to the Singapore Package Insert (PI) or e-mail: [MedInfo.SG@astrazeneca.com](mailto:MedInfo.SG@astrazeneca.com).  
In case of safety concerns, please call +65 6812 4700 or e-mail: [PatientSafety.SG@astrazeneca.com](mailto:PatientSafety.SG@astrazeneca.com).

**Patient name:** \_\_\_\_\_

**Hospital where treated:** \_\_\_\_\_

**Physician name:** \_\_\_\_\_

**Tel. number:** \_\_\_\_\_

**Vaccination date:** \_\_\_\_\_

This document has been approved by Singapore Health Sciences Authority (HSA) on 21-08-2024.



AstraZeneca Singapore Pte Ltd  
10 Kallang Avenue #12-10, Aperia Tower 2, Singapore 339510  
Tel: (+65) 6812 4700, Fax: (+65) 6385 3050